

Function of connexins in the renal circulation

C Wagner¹

¹Physiologisches Institut der Universität Regensburg, Regensburg, Germany

Connexins form intercellular channels that span two plasma membranes and directly couple the cytoplasm of adjacent cells. This morphological contact enables the exchange of ions, second messengers, and metabolites, which act to regulate several biological functions. This review focuses on the significance of connexins in the renal circulation. Cells of the renal vasculature are coupled and express connexins in a vessel and cell-specific pattern. This finding indicates that renal connexins likely play an important role in renal autoregulatory mechanisms (Bayliss effect, tubuloglomerular feedback) and in the control of vasomotor responses. The described coupling of endothelial and vascular smooth muscle cells in the afferent arterioles may also contribute to the communication of neighboring nephrons, called 'nephron coupling.' Furthermore, deletion of the Cx40 and Cx43 genes results in an altered functional behavior of the renin-producing cells, suggesting involvement of these connexin isoforms in the regulation of renin secretion and synthesis. In addition, this review discusses the role of renal connexin expression in the pathogenesis of hypertension or diabetes.

Kidney International (2008) **73**, 547–555; doi:10.1038/sj.ki.5002720; published online 12 December 2007

KEYWORDS: endothelial cells; renal hypertension; renin-angiotensin system; vascular; renal hemodynamics

CONNEXINS: GENERAL STRUCTURE

Connexins are integral membrane proteins that assemble in large transmembrane channels, named connexons or hemichannels. Gap junctions, which can be investigated by electron microscopy, electrophysiological methods, immunohistochemistry, or dye transfer experiments, are composed of two hemichannels (connexons). These connexons are expressed in the plasma membrane of opposing cells and form functional intercellular channels, which directly link the cytoplasm of two adjacent cells. Each connexon consists of six connexin (Cx) protein subunits that enclose a central aqueous pore. The effective pore diameter varies over a range of 0.8–1.6 nm and is determined by the connexin isoform.^{1–3} To date, the connexin gene family comprises of at least 20 different connexin genes in the mouse genome and 21 in the human genome, with significant homology of the respective connexins between human and mouse.^{4–6} Members of the connexin family are distinguished from each other by their predicted molecular weights expressed in kilodaltons (for example, Cx40 is 40 kDa).

The single connexin protein is made up of four-membrane-spanning domains with two conserved extracellular loops that are the essential elements of the docking process, one cytoplasmic loop, and cytoplasmic N- and C-terminal regions (Figure 1), which are the most variant portions of the protein.^{6,7} Connexins can assemble into homomeric connexons when six identical connexins combine to form the hemichannel. Additionally, they can also form heteromeric channels, consisting of two or more connexin variants. Connexons subsequently combine with identical or different connexons and form homo- or heterotypic gap junctions.^{8–10} Heterotypic gap junction channels may be composed of homomeric and/or heteromeric connexons (Figure 1).¹¹ Thus, an unexpected complexity exists in the composition of gap junctions, whereas not all connexins are compatible to form heteromeric or heterotypic channels.^{12,13} The particular combination of connexin isoforms within the channels contributes to the distinct electrical and biochemical properties of the gap junctions.¹⁴ Apart from the isoform composition, channel conductance can be modulated by various factors. The gating properties of gap junctions depend on stimuli such as pH, Ca²⁺, transmembrane and transjunctional voltage,¹⁵ or protein phosphorylation events that lead to modifications in tyrosine, serine, and threonine residues and have been reported to affect intercellular communication.^{16–19} Furthermore, channel conductance can

Correspondence: C Wagner, Physiologisches Institut der Universität Regensburg, Regensburg D-93040, Germany.
E-mail: charlotte.schmid@vkl.uni-regensburg.de

Received 18 June 2007; revised 21 September 2007; accepted 24 October 2007; published online 12 December 2007




Figure 1 | Connexin structure. Schematic model illustrating assembly of connexins into gap junctions. Six connexin subtypes oligomerize into connexons or hemichannels, which combine into homotypic, heterotypic/homomeric, or heterotypic/heteromeric gap junction channels. The connexin protein creates four membrane domains (m1–m4) with two extracellular loops (el 1 + 2), one cytoplasmic loop (cl), and the N and C termini exposed to the cytoplasm.

be regulated by the number of channels, which is controlled by the up- and downregulation of connexin synthesis, assembly, trafficking, and the rate of degradation.^{20,21}

CONNEXINS: SIGNAL TRANSDUCTION

Connexins form gap junctions that allow the diffusional swap of various molecules up to a molecular mass of 1 kDa (for example, ions, second messengers, or metabolites)^{8,19,22,23} and present the molecular basis of direct intercellular signaling. The type and number of molecules that spread from cell to cell through gap junctions in a given time period depends on the diameter, size, and charge of molecules in the pore. There is a common agreement that not only monovalent and divalent ions such as sodium, potassium, calcium, and chloride but also uncharged molecules including glucose can pass through most gap junctions. Apart from calcium, other relevant signaling molecules such as cAMP, cGMP, inositol trisphosphate, and ATP can spread from cell to cell through gap junctions.^{24–27} It has been demonstrated that connexin isoforms can influence the permeability of gap junctions to signaling molecules. For example, gap junctions composed of Cx43 have a considerably higher permeability to AMP, ADP, and mainly ATP compared to Cx32 channels, although adenosine traverses Cx32 gap junction channels at a 12-fold higher rate.²⁷ The permeability of gap junctions is subject to regulation. Thus, calcium seems to modulate the passage of molecules through gap junctions, as high or more focal elevations of intracellular Ca^{2+} act to close gap junction channels,²⁸ whereas lower Ca^{2+} levels exert no effect on gap junctions.

CONNEXINS: GENERAL FUNCTIONS

Except for a few tissues like erythrocytes, thrombocytes, mature skeletal muscle fibers, and spermatozooids, gap junctions are present in all vertebrate cell types.²⁹ However, the expression of certain connexins is specific relative to tissue and cell type. Cell-to-cell channels are involved in the regulation of several physiological and pathophysiological functions such as cell growth,³⁰ glandular secretion³¹ proliferation and differentiation, developmental processes, tumorigenic processes, and regeneration and so on.^{32,33}

Mutations in connexin genes have been associated with a number of human diseases, such as chromosome X-linked Charcot–Marie–Tooth disease,³⁴ which is caused by mutations in Cx32,^{35,36} congenital deafness and skin diseases, which have been linked to mutations in human Cx26,³⁷ and congenital cataracts, which are affected by mutations in Cx50.³⁵

CONNEXINS: SPECIAL FUNCTION

Gap junctions assure the mechanical and electrical coupling of different cell types. Such a role is critical in tissues like the mammalian nervous system, in which gap junctions provide a mode of direct interneuronal communication between adjacent neuronal cells in addition to indirect transmission via chemical synapses,^{38–43} and in the heart, where gap junctions permit the rapid cell-to-cell transfer of electrical impulses and guarantee the coordinated contraction of cardiomyocytes.^{44–49}

Gap junctions are also present in the vascular system. Communication via gap junctions has been demonstrated to

be an important mechanism in the control of vascular function where they are thought to facilitate communication between cells within the blood vessel wall and coordinate the regulation of vascular tone.⁵⁰ Thus, the coupling of vascular smooth muscle cells may be responsible for the synchronous contraction of small blood vessels, whereas junctions between endothelial and vascular smooth muscle cells, the so-called myoendothelial gap junctions (MEGJs), are thought to transmit signals from the endothelium to the underlying smooth muscle cells. Finally, gap junctions between endothelial cells might transport intracellular signals to endothelial cells of upstream arterioles, causing a conducted vasomotor response.⁵¹ The main connexins expressed in vascular cells are Cx37, Cx40, Cx43, and Cx45.^{52,53} Vascular connexins, however, show no homogenous expression pattern between different tissues and even between cell types within the same tissue.

In addition to the broad range of human diseases implicated with mutations in connexin genes, numerous studies have been undertaken in animals deficient in different connexins to determine their physiological function. For example, mice with genetic disruptions in vascular connexins show abnormalities in vascular tone and in propagation of vasomotor responses. Thus, *Cx40* knockout mice exhibit hypertension and show impaired conduction of endothelium-dependent vasodilator responses along arterioles,^{54,55} which indicates a role of Cx40 not only in the control of blood pressure but also in the propagation of vasodilation. Mice with an endothelium-specific deletion of Cx43 circumventing the neonatal lethality of *Cx43* gene disruption are described as normotensive^{54,56} or soft hypotensive.⁵⁷ The absence of Cx45 causes vascular malformations during development and blocks the differentiation of smooth muscle cells, resulting in the narrowing of some vessels and early death.⁵⁸ Observations obtained from these experiments attest to the physiological significance of connexins or, rather, gap junctions in the development and regulation of vascular function.

The renal vasculature does not only regulate the renal blood flow, but is also involved in the control of glomerular responses, thus making the renal vasculature an interesting system for the examination of vascular gap junctions.

CONNEXINS: LOCALIZATION IN THE RENAL VASCULATURE

The distribution of connexins in the vascular system has been investigated in a series of studies that have detected Cx37, Cx40, Cx43, and Cx45 as the major connexins expressed in the vessel walls. Although connexins are not uniformly expressed in all blood vessels and vary between different species, arterial endothelial cells most notably seem to express Cx37 and Cx40, whereas smooth muscle cells express Cx43 and Cx45. In the following section, the localization of connexins will be described explicitly in the vasculature of the kidney (Table 1).

In large renal vessels (for example, renal, interlobar, and arcuate arteries), the connexin isoforms Cx37, Cx40, and

Table 1 | Summarization of the connexin distribution in renal arteries of the mouse

	Large renal arteries		Renal arterioles		
	Endothelium	Media	Endothelium	Media	Renin-producing cells
Cx37	++	+	++	–	+
Cx40	+++	–	+++	–	+++
Cx43	+	+	+	+	–

Cx43 were found in the endothelial layer,^{59–62} although Cx43 expression is weak and irregular. This finding is consistent with previous studies showing Cx43 in the endothelium of rat aorta or carotid artery,^{63,64} but this is in contrast to a recent study in which Cx43 was not found in the mouse aortic endothelium,⁶⁵ indicating substantial species and tissue differences. However, information about connexins in the arterial media of large renal vessels currently available from the literature is insufficient and confusing. Although it is generally described for major arteries that smooth muscle cells express Cx37, Cx40, Cx43, and Cx45,⁶⁰ the exact connexin expression in the media of larger renal arteries is still unclear. Currently, there is immunohistochemical evidence that Cx37 and Cx43 are expressed in the media of large renal arteries.^{61,66}

Much more information is available from the distribution of connexins in renal arterioles. The endothelial cells express Cx37 and Cx40 and weakly express Cx43,^{59–61,67} which is largely consistent with different species and agrees with the pattern of extrarenal arterioles.^{53,54,68,69} In the media of afferent arterioles, connexin subtypes Cx37 and Cx43 are expressed, whereas Cx37 was only found in the preglomerular vessels of mice,⁶¹ but not in the rat kidney.⁷⁰ Investigations of rat and human kidneys show either no expression of Cx43 in the smooth muscle cells of afferent arterioles^{67,70,71} or only weak expression in the media of the afferent arterioles.⁵⁹ However, investigating connexins in the afferent arterioles in more detail reveals an interesting heterogeneous expression pattern of connexins along the arterioles. Although the proximal portions of the afferent arterioles show typical Cx40, Cx37, and also weak Cx43 expression in the endothelium, no gap junctional coupling through these connexin isoforms can be detected in the distal parts of these vessels, in which Cx40 can be found in the smooth muscle cell layer.⁷⁰ Consistent with this result, remarkable Cx40 immunoreactivity was observed not only in the renin-producing cells of the juxtaglomerular apparatus (JGA), considered to be modified smooth muscle cells, but also in extraglomerular and intraglomerular mesangial cells.^{59–62,67} Cx37 is expressed in renin-producing cells as well as in the extraglomerular mesangium and mesangial cells near the vascular pole.⁶¹ In contrast to the afferent arterioles, only Cx43 was detected in the endothelial cells of efferent arterioles and there was no connexin expression in the media, suggesting extensive coupling of the cells in the preglomerular vasculature but not in postglomerular vessels.⁶¹

In summary, cells in the wall of the large renal vessel appear to have a relatively normal connexin expression pattern, suggesting that connexins exert physiological functions, which correspond to their role in the general vascular system. In contrast, connexin expression, especially in the distal part of the afferent arterioles, completely differs from the common pattern by striking extensive but heterogeneous Cx40 expression, which may contribute to the regulation of glomerular function or renal endocrine function.

CXS: ROLE IN RENAL HEMODYNAMICS

The control of renal blood flow is determined by sympathetic nerves, by endothelial function, and by autoregulatory mechanisms. The latter comprise mainly a myogenic component, commonly known as the Bayliss mechanism and the tubuloglomerular feedback (TGF, see below). This myogenic reaction has been demonstrated in a previous study, where preparations of blood-perfused juxtamedullary nephrons abrupt increases in perfusion pressure result in the vasoconstriction of afferent arterioles in a spatially organized manner.⁷² As it is assumed that the myogenic response is activated by a depolarization that opens voltage-dependent channels and leads to an inward Ca^{2+} current, one may hypothesize that the electrotonic diffusion of Ca^{2+} may occur throughout gap junctions connecting the medial layer.⁷³ In fact, several studies in the literature have verified the electrical coupling of smooth muscle cells in afferent arterioles using electrophysiological methods.^{74,75} This assumption was further supported by studies conducted in neuronal arteries, in which the uncoupling of gap junctions by the inhibitor α -glycylrrhentic acid significantly impaired myogenic vasoconstriction.⁷⁶ However, there are no studies detailing the involvement of connexin proteins in mediating the Bayliss effect.

The contractile status of smooth muscle cells can be further regulated by the endothelial release of vasoactive substances, so-called autocooids, which include nitric oxide (NO), prostaglandins, or the endothelium-derived hyperpolarizing factor.^{77,78} It is conceivable that the relaxing action of autocooids on smooth muscle cells is mediated via MEGJs, which couples endothelium cells directly with the media layer. MEGJs have been described in some extrarenal vascular tissues, mostly in smaller arterioles, to be involved in vasotonus control.^{79–81} In renal vessels, Cx40-containing gap junctions are present in endothelial and renin-producing cells,⁶⁷ although no evidence for MEGJs among normal smooth muscle and endothelial cells exists. Although NO or PGI_2 presumably diffuse from the endothelium to the smooth muscle cells, subsequent relaxation is not affected by gap junction blockers.⁸² Endothelium-derived hyperpolarizing factor activity, in contrast, seems to be dependent on gap junctional coupling.⁸³ Information about certain connexins involved in mediating the vasomotor responses is limited and only available from studies conducted on extrarenal tissues such as rat mesenteric and basilar arteries, in which MEGJs are described to be composed of Cx37 and

Cx40.^{84,85} Furthermore, in human subcutaneous arterioles, endothelium-derived hyperpolarizing factor-induced vasodilatation was shown to be dependent on Cx43 using the mimetic peptide ⁴⁰Gap26.⁶⁹

In the microcirculatory system, the vasomotor responses can spread along the vessel wall to facilitate dilatation or constriction of upstream arterioles according to blood flow requirements. The propagation of vasodilatation or vasoconstriction is linked to an electric conduction of a hyperpolarizing or depolarizing current, which is known to flow through gap junctions⁸⁶ and couples cells of either the endothelium or the smooth muscle of arterioles. Conduction of depolarizing currents has already been shown in renal afferent arterioles in rats, where the microapplication of KCl causes the spread of a local vasoconstriction over a distance up to 1500 μm ^{87,88} to produce a sufficient increase in afferent arteriolar resistance. The conduction of vasodilatation has also been shown to be dependent on intercellular coupling. In skeletal muscle arterioles of mice deficient for Cx40, conduction of vasodilatation was markedly impaired,^{54,89} suggesting that the vasodilatory response requires cell coupling via Cx40. The question remains whether conduction of the vasomotor responses is conducted along the endothelium or along the media of the arterioles. To explore this issue, investigations in mice where the continuity either in endothelium or smooth muscle cells was interrupted show that vasodilatation is conducted along the endothelium, whereas vasoconstriction is conducted in the media.^{79,90} This assumption fits well with the attenuated conductance of vasodilatation in mice lacking Cx40, which is known to be the predominant endothelial connexin protein.⁵⁴ The role of specific connexins in the regulation of the vasomotor responses in the kidney is currently unknown and requires further investigation.

CXS: ROLE IN TGF SIGNALING

The glomerular filtration rate of the kidney is controlled via the TGF mechanism. For this sophisticated feedback loop, a specialized site of the nephron (the macula densa) contacts its own afferent arteriole in the JGA. The macula densa cells sense increasing sodium chloride concentrations in the tubule lumen and send vasoconstrictor signals to smooth muscle cells inside the wall of the afferent arteriole to regulate the preglomerular diameter and thereby decrease the glomerular filtration of the respective nephron.^{91,92} This process requires transfer of the vasoconstrictor signal from macula densa cells via the strategically positioned extraglomerular mesangial cells to the smooth muscle cells in the afferent arteriole.^{93–95} As shown in a previous study, cells of the JGA, except those of the macula densa,⁹⁶ are coupled via gap junctions^{97,98} to establish a pathway for the transmission of the vasoconstrictor signal. Indeed, the importance of gap junctions within the JGA and their role in TGF was recently presented in a study where damaging mesangial cells with a Thy 1-1 antibody and the disruption of gap junctions by heptanol led to the elimination of TGF.⁹⁹ Heptanol treatment does not compromise the function of JGA cells. A study by

Ren *et al.*⁹⁹ was unable to elucidate whether TGF requires coupling between the mesangial cells, an intercellular link between mesangial cells and smooth muscle cells, or a continuous connection between these different cell types.¹⁰⁰ Information about the connexin subtypes that determine the TGF has not been elucidated, although numerous findings suggest that Cx40 is the main junctional protein in the JGA.^{60,61,67,70} In agreement with these investigations, we recently observed that the genetic disruption of the *Cx40* gene in mice is associated with increased interstitial spaces and the failure of intercellular coupling between extraglomerular mesangial cells.¹⁰¹ To gain greater insight into the function of certain connexins such as Cx40 in TGF signaling, additional experiments will be necessary to study whether the absence of connexin proteins allows the transmission of macula densa signals to the afferent arteriole.

There is accumulating evidence that ATP is released from macula densa cells in response to increasing sodium chloride concentration in the tubule lumen.^{102–104} As there is no cell coupling between macula densa cells and mesangial cells, it appears that macula densa cells release ATP at their basolateral membrane probably through connexin hemichannels, which may provide a pathway for the release of paracrine signals.^{105–107} The binding of ATP to P₂X or P₂Y receptors, which are abundantly expressed in cells of the JGA,^{108–111} triggers a calcium wave that spreads from the macula densa via the extraglomerular mesangium toward the afferent arterioles.⁹⁵ Interestingly, the uncoupling of gap junctions by gap junction uncoupling agents, α -glycyrrhethinic acid or heptanol, abrogates increases in intracellular calcium and the spreading of the calcium wave,^{95,112} suggesting an essential role of cell coupling in the TGF mechanism.

In summary, although there is accumulating evidence that gap junctions play a role in mediating TGF signaling, detailed information about the localization of participating gap junctions in the JGA or involved connexin subtypes is still lacking.

More studies are necessary to define the exact role of connexins in the regulation of the TGF mechanism.

CXS: ROLE FOR NEPHRON COUPLING

The TGF controls regular oscillations of hydrostatic pressure in the glomerular capillaries. The capillary pressure of two neighboring glomeruli often oscillates synchronously.^{87,113,114} As experimentation of the TGF response at a defined glomerulus also leads to an almost simultaneous TGF response in a neighboring glomerulus, it is obvious that two neighboring glomeruli communicate with each other, a phenomenon that has been termed ‘nephron coupling.’ The structural basis for this nephron coupling is bi- or trifurcated endings of afferent arterioles. The nephron coupling mechanism involves retrograde signal propagation from one vascular pole to the branching sites followed by orthograde propagation of the signal into the other branches.⁷² The signal itself must influence the tone of the

afferent arterioles, in particular, at the entrance into the glomerulus. The signal is conducted along the walls of the afferent arterioles through gap junctions. In principle, signal conduction could also occur along the endothelial or the smooth muscle cell layers. From the available data, it is not yet possible to attribute nephron coupling to endothelial or smooth muscle cell gap junctions. Knockout mice should provide a powerful tool that could aid in addressing this issue. For example, endothelial conduction could be expected to be impaired in *Cx40*-deficient mice, whereas mice with a vsmc-specific disruption of the respective connexin genes should show altered conduction along the media layer. The nature of the propagated signal will likely be characterized in future experiments. The simplest explanation is that changes in membrane potential are propagated from one afferent arteriole to another.⁸⁷ As the smooth muscle cells of afferent arterioles contain calcium channels,^{115,116} changes in the membrane potential would be rapidly transformed into changes in vessel tone. In this context, Wagner *et al.*⁸⁷ have demonstrated that vasoconstriction induced at the distal end of a certain afferent arteriole by KCl application was conducted in the cortical radial artery and adjacent afferent arterioles in the rat kidney. Interestingly, the conduction of these vasomotor responses was much greater in spontaneous hypertensive rats, indicating strengthening of internephron coupling in the hypertensive state. One may hypothesize that the known enhanced Cx40 and Cx43 expression during hypertension (see section below) may improve the intercellular coupling within the vessel wall, thus causing the intensified conduction of neighboring nephrons in spontaneous hypertensive rats. However, this issue is still unclear and requires further investigation to determine whether the hypertension-induced elevation of connexins primarily affects endothelium coupling or junctional communication between smooth muscle cells.

CXS: REGULATION OF RENIN SYNTHESIS AND SECRETION

The JGA represents a multilayer morphological structure, which comprises of several cell types in a very small kidney compartment. It has been known that these cells are strongly coupled via gap junctions. Morphological evidence has been corroborated by functional data indicating strong electrical coupling of renin-secreting cells.^{74,75,117} Within the JGA, renin-producing cells form numerous gap junctions between renin-producing cells themselves as well as between renin-producing cells and the adjacent endothelial or extraglomerular mesangial cells.^{97,118} The predominant isoform of connexins in renin-producing cells is Cx40 (Figure 2), which is expressed in high density in these cells.^{60,63,66} The expression pattern of Cx40 is remarkable. Within the vasculature, Cx40 is expressed almost exclusively in the endothelium. For renin-producing cells that are known to replace vascular smooth muscle cells in the media of distal portions of the afferent arterioles, one would expect that they do not express connexins, which are typical for the endothelium, but specifically connexin subtypes including




Figure 2 | Cx40 expression in the afferent arteriole.

Immunofluorescence analysis of cryostat sections of mouse kidneys shows Cx40 expression in the endothelium of the afferent arteriole (indicated by arrows) and in renin-producing cells (rpc). Green staining for renin; red staining for Cx40.

Cx43, which are normally found in vascular smooth muscle cells. Interestingly, juxtaglomerular renin-producing cells do not express Cx43.^{59,67,70} Although the functional relevance of the striking cell coupling in the JGA has not been extensively studied, one may hypothesize that Cx40-mediated contact in renin-producing cells with either the endothelium or mesangial cells provides an ideal pathway for signals involved in controlling the activity of the renin system. Synthesis and secretion of renin are essentially controlled by circulating angiotensin II levels, the sodium chloride concentration passing the macula densa, and the classic negative feedback loops such as the hydrostatic pressure within the vessel. Most recently, the role of Cx40 gap junctions in the transmission of inhibitory signals has been addressed in studies using the *Cx40* knockout model.^{62,119} The classic approach to investigate the influence of blood pressure on renin secretion is the experimental unilateral renal artery stenosis, which led to an increased renin secretion in the hypoperfused kidney causing hypertension and decreased secretion in the contralateral intact kidney. Performing this maneuver in *Cx40*-deficient mice did not lead to an increase in blood pressure and plasma renin concentration and did not promote the expected increased renin secretion in the hypoperfused kidney.^{62,119}

Investigations of the inhibitory action of Ang II on the renin system led to similar results. This has been shown by *in vivo* experiments in which diminished circulating Ang II levels resulting from ACE inhibition led to a clearly reduced stimulation of renin secretion and renin synthesis in *Cx40*^{-/-} mice.^{62,119} Finally, it could be shown that the deficiency of *Cx40* interrupts the inhibitory macula densa signaling of renin-producing cells. Thus, the application of loop diuretics normally stimulates the activity of the renin system in wild-type mice, whereas this stimulating effect is markedly attenuated in *Cx40* knockout mice.¹⁰¹ These findings indicate that the classic feedback loops, which involve intrarenal blood pressure, Ang II, or salt load rely on cell-to-cell transmission

of respective inhibitory signals of renin-producing cells and are disturbed in the absence of Cx40.

Interestingly, mice where *Cx43* (normally expressed in rare endothelial cells and smooth muscle cells in afferent arterioles,^{61,66,67,70,119} but not in renin-producing cells) is replaced by *Cx32* (*Cx43ki32*) displayed markedly reduced renin secretion and failure of hypertension in the model of unilateral renal stenosis.¹²⁰ These findings indicate that intercellular communication via Cx40 expressed by renin-producing cells, and also via Cx43 located in endothelial cells, is required for control of the renin system. How the disruption of *Cx40* or *Cx43* genes can exert such characteristic changes in the activity of the renin system is yet unknown and requires further investigation. One could speculate that the overlapping defects in the control of the renin system seen by replacement of *Cx43* or by the deletion of *Cx40* could support a functional link of Cx40 and Cx43 signaling mechanism. As both isoforms do not appear to be compatible to form heterotypic channels,¹²¹ one could imagine that due to the spatial vicinity of the cells within the JGA, transmission of humoral factors by paracrine signaling may also play a certain role for the control of the renin system.

CXS: MODULATION OF THEIR EXPRESSION IN THE KIDNEY

Connexins have been described in numerous studies to be involved in the development of vascular pathophysiology in that they modulate their expression levels.^{50,54,122,123}

For example, diabetes induction by intraperitoneal injections of streptozotocin causes a modified expression of Cx40 and Cx43 in the renal vasculature of C57Bl/6 mice. In controls, Cx40 is expressed in the endothelium of the afferent arterioles and appears in the media only in the terminal part of these vessels. However, Cx43 was detected in endothelial cells of the efferent arterioles as described above. In diabetes, increased Cx40 expression was observed in vascular smooth muscle cells along the afferent arterioles and inside the glomerulus. However, Cx43 staining in the endothelium along the efferent arteriole decreased considerably, indicating that during diabetes the intercellular communication via gap junction is enhanced in the preglomerular vasculature, but concurrently abrogated in the postglomerular parts.^{61,124} These observations are supported by an *in vitro* study that demonstrates the downregulation of Cx43 in mesangial cells exposed to a high glucose medium.¹²⁵ In addition to the altered connexin expression in diabetes, the availability of NO increases with this disease. Zhang *et al.*¹²⁴ investigated the possible role of NO on the changes in connexin expression. In a very elegant experimental strategy, connexin expression was studied in mice with overexpression or deletion of endothelial NO synthase activity. Interestingly, mice overexpressing endothelial NO synthase showed changes in Cx40 and Cx43 expressions similar to the modified connexin expression in wild-type mice during diabetes. Basal Cx40 and Cx43 expressions in endothelial NO synthase knockout mice do not differ from wild-type mice, and

the induction of diabetes produces no alteration of connexins either in preglomerular or postglomerular parts of the renal vasculature. This leads to the assumption that NO, which is regulated by endothelial NO synthase activity, may contribute to the changes in connexin expression during diabetes. This is in contrast to a subsequent *in vitro* study that also investigated the involvement of NO in the control of gap junctions and Cx43 expression, but exhibited NO as a potent Cx43 stimulator.¹²⁶

Modifications of connexins in the kidney have been further described in studies interested in the pathogenesis of hypertension. One of these studies investigated the effects of renovascular hypertension induced by unilateral renal artery clipping. As shown by molecular biological analysis, experimentally induced high blood pressure leads to an increase in Cx40 mRNA and protein in the clipped and non-clipped kidneys, whereas Cx43 was only increased in the non-clipped kidney. Immunohistochemical analysis revealed that the elevation of Cx40 protein was due to its enhanced expression in renin-producing cells in the wall of afferent arterioles.⁶⁷ Similarly, Cx43 expression is also affected by hypertension in the aortic wall. Independent of the generating mechanism for hypertension, Cx43 expression is markedly increased in different hypertensive models such as the 2K1C, Doca-salt, or L-NAME (*N*_ω-nitro-L-arginine methyl ester hydrochloride) model, which suggests that mechanical forces induced by blood pressure elevation mainly regulate Cx43 expression.^{67,127}

In summary, connexins appear to play an important role in the development of vascular diseases. Apparently, the modified cell-to-cell communication between vascular cells disturbs the regulation of normal vascular function causing the pathogenesis of diabetic nephropathy or hypertension, for example.

CXS IN RENAL VASCULATURE: CONCLUSIONS

As there is only limited data to document the function of connexins in the renal vascular system, previous sections have provided support for more possible roles of connexins in the regulation of renal vascular hemodynamics, TGF signaling, nephron coupling, or the renin system. One could infer that if connexins play such a critical role in maintaining renal circulatory function, they are potential molecular targets for the pathogenesis of renal vascular disease. However, it must be stressed that there is presently too little information regarding a direct role of connexins in vascular disease. Interestingly, a polymorphism within the Cx40 promoter has been shown to be associated with increased risk of hypertension in men.¹²⁸ Thus, it remains to be elucidated whether changes of renal connexin expression account for genesis of vascular disease in humans. Given this background, it seems to be a very interesting field for additional research.

ACKNOWLEDGMENTS

The author thanks Dr Armin Kurtz for scientific discussions. Lisa Kurtz provided the immunohistochemical image. C Wagner was supported by a grant from Deutsche Forschungsgemeinschaft (SFB699/B2).

REFERENCES

- Veenstra RD. Ion permeation through connexin gap junction channels: effects on conductance and selectivity. In: Peracchia C (ed). *Gap Junctions: Molecular Basis of Cellular Communication in Health and Disease. Current Topics in Membranes*, vol. 49. Academic Press: New York, 2000 pp 95–129.
- Gong XQ, Nicholson BJ. Size selectivity between gap junction channels composed of different connexins. *Cell Commun Adhes* 2001; **8**: 187–192.
- Weber PA, Chang H-C, Spaeth KE *et al*. The permeability of gap junction channels to probes of different size is dependent on connexin composition and permanent-pore affinities. *Biophys J* 2004; **87**: 958–973.
- Sohl G, Willecke K. An update on connexin genes and their nomenclature in mouse and man. *Cell Commun Adhes* 2003; **10**: 173–180.
- Evans WH, Martin PE. Gap junctions: structure and function. *Mol Membr Biol* 2002; **19**: 121–136.
- Soehl G, Maxeiner S, Willecke K. Expression and functions of neuronal gap junctions. *Nat Rev* 2005; **6**: 192–200.
- Soehl G, Willecke K. Gap junctions and the connexin protein family. *Cardiovasc Res* 2004; **62**: 288–292.
- Goodenough DA, Goliger JA, Paul DL. Connexins, connexons, and intercellular communication. *Annu Rev Biochem* 1996; **65**: 474–502.
- White TW, Paul DL. Genetic diseases and knockouts reveal diverse connexin functions. *Annu Rev Biochem* 1999; **65**: 474–502.
- Willecke K, Eiberger J, Degen J *et al*. Structural and functional diversity of connexin genes in the mouse and human genome. *Biol Chem* 2002; **383**: 725–737.
- Cottrell GT, Burt JM. Heterotypic gap junction channel formation between heteromeric and homomeric Cx40 and Cx43 connexons. *Am J Physiol Cell Physiol* 2001; **281**: C1559–C1567.
- Jiang JX, Goodenough DA. Heteromeric connexons in lens gap junction channels. *Proc Natl Acad Sci USA* 1996; **93**: 1287–1291.
- Rackauskas M, Verselis VK, Bukauskas FF. The permeability of homotypic and heterotypic gap junction channels formed of cardiac connexins, mCx30.2, Cx40, Cx43 and Cx45. *Am J Physiol Heart Circ Physiol* 2007; **293**: H1729–H1736.
- Harris AL. Emerging issues of connexin channels: biophysics fills the gap. *Q Rev Biophys* 2001; **34**: 325–472.
- Sosinsky GE, Nicholson BJ. Structural organization of gap junction channels. *Biochim Biophys Acta* 2005; **1711**: 99–125.
- Lampe PD, Lau AF. Regulation of gap junctions by phosphorylation of connexins. *Arch Biochem Biophys* 2000; **384**: 205–215.
- Lampe PD, Lau AF. The effects of connexin phosphorylation on gap junctional communication. *Int J Biochem Cell Biol* 2004; **36**: 1171–1186.
- Moreno AP, Lau AF. Gap junction channel gating modulated through protein phosphorylation. *Prog Biophys Mol Biol* 2007 (in press).
- Bruzzone R, White TW, Paul DL. Connections with connexins: the molecular basis of direct intercellular signalling. *Eur J Biochem* 1996; **238**: 1–27.
- Segretain D, Falk MM. Regulation of connexin biosynthesis, assembly, gap junction formation, and removal. *Biochim Biophys Acta* 2004; **1662**: 3–21.
- Brink PR. Gap junctions in vascular smooth muscle. *Acta Physiol Scand* 1998; **164**: 349–356.
- Kumar NM, Gilula NB. The gap junction communication channel. *Cell* 1996; **84**: 381–388.
- Saez JC, Berthoud VM, Branes MC *et al*. Plasma membrane channels formed by connexins: their regulation and functions. *Physiol Rev* 2003; **83**: 1359–1400.
- Lawrence TS, Beers WH, Gilula NB. Transmission of hormonal stimulation by cell-to-cell communication. *Nature* 1978; **272**: 501–506.
- Saez JC, Connor JA, Spray DC *et al*. Hepatocyte gap junctions are permeable to the second messenger, inositol 1,4,5-trisphosphate, and to calcium ions. *Proc Natl Acad Sci* 1989; **86**: 2708–2712.
- Sanderson M. Intercellular calcium waves mediated by inositol triphosphate. *Ciba Found Symp* 1995; **188**: 175–194.
- Goldberg GS, Moreno AP, Lampe PD. Gap junctions between cells expressing connexin 43 or 32 show inverse permselectivity to adenosine and ATP. *J Biol Chem* 2002; **278**: 46533–46540.
- Dakin K, Zhao YR, Li W-H. LAMP, a new imaging assay of gap junctional communication unveils that Ca²⁺ influx inhibits cell coupling. *Nat Methods* 2005; **2**: 55–62.
- Willecke K, Eiberger J, Degen J *et al*. Structural and functional diversity of connexin genes in the mouse and human genome. *Biol Chem* 2002; **383**: 725–737.
- Loewenstein WR, Rose B. The cell-cell channel in the control of growth. *Semin Cell Biol* 1992; **3**: 59–79.

31. Meda P, Chanson M, Pepper M *et al.* *In vivo* modulation of connexin 43 gene expression and junctional coupling of pancreatic β -cells. *Exp Cell Res* 1991; **192**: 469–480.
32. Wei CJ, Xu X, Lo CW. Connexins and cell signaling in development and disease. *Annu Rev Cell Dev Biol* 2004; **20**: 811–838.
33. Jiang JX, Gu S. Gap junction- and hemichannel-independent actions of connexins. *Biochim Biophys Acta* 2005; **1711**: 208–214.
34. Bergoffen J, Scherer SS, Wang S *et al.* Connexin mutations in X-linked Charcot-Marie-Tooth disease. *Science* 1993; **262**: 2039–2042.
35. Krutovskikh V, Yamasaki H. Connexin gene mutations in human genetic diseases. *Mutat Res* 2000; **462**: 197–207.
36. Zhou L, Griffin JW. Demyelinating neuropathies. *Curr Opin Neurol* 2003; **16**: 307–313.
37. Liu XZ, Walsh J, Mburu P *et al.* Mutations in the myosin VIIA gene cause non-syndromic recessive deafness. *Nat Genet* 1997; **16**: 188–190.
38. Bennett M, Zukin RS. Electrical coupling and neuronal synchronization in the mammalian brain. *Neuron* 2004; **41**: 495–511.
39. Nagy JI, Dudek JE, Rash JE. Update on connexins and gap junctions in neurons and glia in the mammalian nervous system. *Brain Res Rev* 2004; **47**: 191–215.
40. Montoro RJ, Yuste R. Gap junctions in developing neocortex: a review. *Brain Res Rev* 2004; **47**: 216–226.
41. Buzsaki G, Chrobak JJ. Temporal structure in spatially organized neuronal ensembles: a role for interneuronal networks. *Curr Opin Neurobiol* 1995; **5**: 504–510.
42. Fricker D, Miles R. Interneurons, spike timing, and perception. *Neuron* 2001; **32**: 771–774.
43. Theis M, Söhl J, Eiberger K *et al.* Emerging complexities in identity and function of glial connexins. *Trends Neurosci* 2005; **28**: 188–195.
44. De Maziere A, Analbers L, Jongsma HJ *et al.* Immunoelectron microscopic visualization of the gap junction protein connexin 40 in the mammalian heart. *Eur J Morphol* 1993; **31**: 51–54.
45. Gros D, Jarry-Guichard T, Ten Velde I *et al.* Restricted distribution of connexin 40, a gap junctional protein in mammalian heart. *Circ Res* 1994; **74**: 839–851.
46. Beyer EC, Paul DL, Goodenough DA. Connexin 43: a protein from rat heart homologous to the gap junction protein from liver. *J Cell Biol* 1987; **105**: 2621–2629.
47. Gros DB, Jongsma HJ. Connexins in the mammalian heart function. *Bioessays* 1996; **18**: 719–730.
48. Coppen SR, Kodema I, Boyett MR *et al.* Connexin 45, a major connexin of the rabbit sinoatrial node, is co-expressed with connexin 43 in a restricted zone at the nodal-crista terminalis border. *J Histochem Cytochem* 1999; **47**: 907–918.
49. Coppen SR, Severs NJ, Gourdie RG. Connexin 45 (alpha 6) expression delineates an extended conduction system in the embryonic and mature rodent heart. *Dev Genet* 1999; **24**: 82–90.
50. Christ GJ, Spray DC, el-Sabban M *et al.* Gap junctions in vascular tissues. Evaluating the role of intercellular communication in the modulation of vasomotor tone. *Circ Res* 1996; **79**: 631–646.
51. Segal SS, Duling BR. Propagation of vasodilation in resistance vessels of the hamster: development and review of a working hypothesis. *Circ Res* 1987; **61**: 1120–1125.
52. Dhein S. Gap junction channels in the cardiovascular system: pharmacological and physiological modulation. *Trends Pharmacol Sci* 1998; **19**: 229–241.
53. Gustafsson F, Mikkelsen HB, Arensbak B *et al.* Expression of connexin 37, 40 and 43 in rat mesenteric arterioles and resistance arteries. *Histochem Cell Biol* 2003; **119**: 139–148.
54. de Wit C, Ross F, Bolz SS *et al.* Impaired conduction of vasodilation along arterioles in connexin 40-deficient mice. *Circ Res* 2000; **86**: 649–655.
55. de Wit C, Roos F, Bolz SS *et al.* Lack of connexin 40 is associated with hypertension and irregular arteriolar vasomotion. *Physiol Genomics* 2003; **13**: 169–177.
56. Theis M, de Wit C, Schlaefer TM *et al.* Endothelium-specific replacement of the connexin 43 coding region by a lacZ reporter gene. *Genesis* 2001; **29**: 1–13.
57. Liao Y, Day KH, Damon DN *et al.* Endothelial cell-specific knockout of connexin 43 causes hypotension and bradycardia in mice. *Proc Natl Acad Sci USA* 2001; **98**: 9989–9994.
58. Kruger O, Plum A, Kim JS *et al.* Defective vascular development in connexin 45-deficient mice. *Development* 2000; **127**: 4179–4193.
59. Hwang Seul K, Beyer EC. Heterogenous localization of connexin 40 in the renal vasculature. *Microvasc Res* 2000; **59**: 140–148.
60. Haefliger JA, Nicod P, Meda P. Contribution of connexins to the function of the vascular wall. *Cardiovasc Res* 2004; **62**: 345–356.
61. Zhang J, Hill CE. Differential connexin expression in preglomerular and postglomerular vasculature: accentuation during diabetes. *Kidney Int* 2005; **68**: 1171–1185.
62. Wagner C, de Wit C, Kurtz L *et al.* Connexin 40 is essential for the pressure control of renin synthesis and secretion. *Circ Res* 2007; **100**: 556–563.
63. Yeh HI, Lupu F, Dupont E *et al.* Upregulation of connexin 43 gap junctions between smooth muscle cells after balloon catheter injury in the rat carotid artery. *Arterioscler Thromb Vasc Biol* 1997; **17**: 3174–3184.
64. Hong T, Hill CE. Restricted expression of the gap junctional protein connexin 43 in the arterial system of the rat. *J Anat* 1998; **192**: 583–593.
65. Simon AM, McWhorter AR, Chen H *et al.* Decreased intercellular communication and connexin expression in mouse aortic endothelium during lipopolysaccharide-induced inflammation. *J Vasc Res* 2004; **41**: 323–333.
66. Barajas L, Liu L, Tucker M. Localization of connexin 43 in the rat kidney. *Kidney Int* 1994; **46**: 621–626.
67. Haefliger JA, Demotz S, Braissant O *et al.* Connexins 40 and 43 are differentially regulated within the kidneys of rats with renovascular hypertension. *Kidney Int* 2001; **60**: 190–201.
68. Looft-Wilson RC, Payne GW, Segall SS. Connexin expression and conducted vasodilation along arteriolar endothelium in mouse skeletal muscle. *J Appl Physiol* 2004; **97**: 1152–1158.
69. Lang NN, Luksha L, Mewby DE *et al.* Connexin 43 mediates endothelium-derived hyperpolarizing factor-induced vasodilation in subcutaneous resistance arteries from healthy pregnant women. *Am J Physiol Heart Circ Physiol* 2007; **292**: H1026–H1032.
70. Arensbak B, Mikkelsen HB, Gustafsson F *et al.* Expression of connexin 37, 40 and 43 mRNA and protein in renal preglomerular arterioles. *Histochem Cell Biol* 2001; **115**: 479–487.
71. Hillis GS, Duthie LA, Mlynski R *et al.* The expression of connexin 43 in human kidney and cultured renal cells. *Nephron* 1997; **75**: 458–463.
72. Casellas D, Bouriquet N, Moore LC. Branching patterns and autoregulatory responses of juxtamedullary afferent arterioles. *Am J Physiol* 1997; **272**: F416–F421.
73. Rummery NM, Hill CE. Vascular gap junctions and implications for hypertension. *Clin Exp Pharmacol Physiol* 2004; **31**: 659–667.
74. Buhle CP, Hackenthal E, Helmchen U *et al.* The hydronephrotic kidney of the mouse as a tool for intravital microscopy and *in vitro* electrophysiological studies of renin-containing cells. *Lab Invest* 1986; **54**: 462–472.
75. Russ U, Rauch U, Quast U. Pharmacological evidence for a K_{ATP} channel in renin-secreting cells from rat kidney. *J Physiol* 1999; **517**: 781–790.
76. Lagaud G, Karicheti V, Knot HJ *et al.* Inhibitors of gap junctions attenuate myogenic tone in cerebral arteries. *Am J Physiol Heart Circ Physiol* 2002; **283**: H2177–H2186.
77. Taylor SG, Weston AH. Endothelium derived hyperpolarizing factor: a new endogenous inhibitor from the vascular endothelium. *Trends Pharmacol Sci* 1988; **9**: 273–274.
78. Busse R, Fleming I. Regulation of endothelium-derived vasoactive autacoid production by hemodynamic forces. *Trends Pharmacol Sci* 2003; **24**: 24–29.
79. Emerson GG, Segal SS. Electrical coupling between endothelial cells and smooth muscle cells in hamster feed arteries: role in vasomotor control. *Circ Res* 2000; **87**: 474–479.
80. Yashiro Y, Duling BR. Integrated Ca^{2+} signalling between smooth muscle and endothelium of resistance vessels. *Circ Res* 2000; **87**: 1048–1054.
81. Sandow SL, Tare M, Coleman HA *et al.* Involvement of myoendothelial gap junctions in the actions of endothelium derived hyperpolarizing factor. *Circ Res* 2002; **90**: 1108–1113.
82. Sandow SL, Bramich NJ, Bandi HP *et al.* Structure, function, and endothelium derived hyperpolarizing factor in the caudal artery of the SHR and WKY rat. *Arterioscler Thromb Vasc Biol* 2003; **2003**: 822–828.
83. Evans WH, Boitano S. Connexin mimetic peptides. Specific inhibitors of gap junctional intercellular communication. *Biochem Soc Trans* 2001; **29**: 606–612.
84. Sandow SL, Neylon CB, Chen MX *et al.* Spatial separation of endothelial small- and intermediate-conductance calcium-activated potassium channels (K_{Ca}) and connexins: possible relationship to vasodilator function? *J Anat* 2006; **209**: 689–698.
85. Haddock RE, Grayson TH, Brackenburg TD *et al.* Endothelial coordination of cerebral vasomotion via myoendothelial gap junctions containing connexin 37 and 40. *Am J Physiol Heart Circ Physiol* 2006; **291**: H2047–H2056.
86. Segal S, Duling BR. Conduction of vasomotor responses in arterioles. A role for cell-to-cell coupling? *Am J Physiol* 1989; **256**: H838–H845.

87. Wagner AJ, Holstein-Rathlou NH, Marsh DJ. Internephron coupling by conducted vasomotor responses in normotensive a spontaneously hypertensive rats. *Am J Physiol* 1997; **272**(3 Part 2): F372–F379.
88. Gustafsson F, Holstein-Rathlou N-H. Conducted vasomotor responses in arterioles: characteristics, mechanisms and physiological significance. *Acta Physiol Scand* 1999; **167**: 11–21.
89. Figueroa XF, Paul DL, Simon AM. Central role of connexin 40 in the propagation of electrically activated vasodilation in mouse cremasteric arterioles *in vivo*. *Circ Res* 2003; **92**: 793–800.
90. Bartlett IS, Segal SS. Resolution of smooth muscle and endothelial pathways for conduction along hamster cheek pouch arterioles. *Am J Physiol Heart Circ Physiol* 2000; **278**: H604–H612.
91. Briggs JP, Schnermann J. The tubuloglomerular feedback mechanism: functional and biochemical aspects. *Annu Rev Physiol* 1987; **49**: 251–273.
92. Navar LG, Inscho EW, Majid DSA *et al*. Paracrine regulation of the renal microcirculation. *Physiol Rev* 1996; **76**: 425–536.
93. Moore LC, Casellas D, Persson AE *et al*. Renal hemodynamic regulation by the renin secreting segment of the afferent arteriole. *Kidney Int Suppl* 1990; **30**: S65–S68.
94. Peti-Peterdi J, Morishima S, Bell PD *et al*. Two-photon excitation fluorescence imaging of the living juxtaglomerular apparatus. *Am J Physiol Renal Physiol* 2002; **283**: F197–F201.
95. Peti-Peterdi J. Calcium wave of tubuloglomerular feedback. *Am J Physiol Renal Physiol* 2006; **291**: 473–480.
96. Bell PD, Lapointe JY, Peti-Peterdi J. Macula densa cell signaling. *Ann Rev Physiol* 2003; **65**: 481–500.
97. Taugner P, Buhle CP, Nobiling R. Ultrastructural changes associated with renin secretion from juxtaglomerular apparatus of mice. *Cell Tissue Res* 1984; **237**: 459–472.
98. Goligorsky MS, Iijima K, Krivneko Y *et al*. Role of mesangial cells in macula densa to the afferent arteriole information transfer. *Clin Exp Pharmacol Physiol* 1997; **24**: 527–531.
99. Ren Y, Carretero OA, Garvin JL. Role of mesangial cells and gap junctions in the tubuloglomerular feedback. *Kidney Int* 2002; **62**: 525–531.
100. Sabirov RZ, Okada Y. ATP-conducting maxi-anion channel: a new player in stress-sensory transduction. *Jpn J Physiol* 2004; **54**: 7–14.
101. Kurtz L, Schweda F, de Wit C *et al*. Lack of connexin 40 causes displacement of renin-producing cells from afferent arterioles to the extraglomerular mesangium. *J Am Soc Nephrol* 2007; **18**: 1103–1111.
102. Castrop H. Mediators of tubuloglomerular feedback regulation of glomerular filtration: ATP and adenosine. *Acta Physiol* 2007; **189**: 3–14.
103. Ren Y, Garvin JL, Liu R *et al*. Role of macula densa adenosine triphosphate (ATP) in tubuloglomerular feedback. *Kidney Int* 2004; **66**: 1479–1485.
104. Sabirov RZ, Sheiko T, Liu H *et al*. Genetic demonstration that the plasma membrane maxi-anion channel and voltage-dependent anion channels are unrelated proteins. *J Biol Chem* 2006; **281**: 1897–1904.
105. Cortina ML, Lin JH, Alves-Rodrigues A *et al*. Connexins regulate calcium signaling by controlling ATP release. *Proc Natl Acad Sci USA* 1998; **95**: 15735–15740.
106. Yao J, Suwa M, Li B *et al*. ATP dependent mechanism for coordination of intercellular Ca²⁺ signaling and renin secretion in rat juxtaglomerular cells. *Circ Res* 2003; **93**: 338–345.
107. Evans WH, De Vuyst E, Leybaert L. The gap junction cellular internet: connexin hemichannels enter the signalling limelight. *Biochem J* 2006; **397**: 1–14.
108. Inscho EW. Renal microvascular effects of P2 receptor stimulation. *Clin Exp Pharmacol Physiol* 2001; **28**: 332–339.
109. Inscho EW, Cook AK, Imig JD *et al*. Physiological role of P2X1 receptors in renal microvascular autoregulatory behaviour. *J Clin Invest* 2003; **112**: 1895–1905.
110. Nishiyama A, Rahman M, Inscho EW. Role of interstitial ATP and adenosine in the regulation of renal hemodynamics and microvascular function. *Hypertens Res* 2004; **27**: 791–804.
111. Schwiebert EM. ATP release mechanism ATP receptors and purinergic signalling along the nephron. *Clin Exp Pharmacol Physiol* 2001; **28**: 340–350.
112. Yao J, Morioka T, Bing L *et al*. Coordination of mesangial cell contraction by gap junction-mediated intercellular Ca²⁺ wave. *J Am Soc Nephrol* 2002; **13**: 2018–2026.
113. Yip KP, Holstein-Rathlou NH, Marsh DJ. Dynamics of TGF-initiated nephron–nephron interactions in normotensive rats and SHR. *Am J Physiol* 1992; **262**(6 Pt 2): F980–F988.
114. Holstein-Rathlou NH, Marsh DJ. Oscillations of tubular pressure, flow, and distal chloride concentration in rats. *Am J Physiol* 1989; **256**(6 Part 2): F1007–F1014.
115. Hansen PB, Jensen BL, Andreassen D *et al*. Differential expression of T- and L-type voltage-dependent calcium channels in renal resistance vessels. *Circ Res* 2001; **89**: 630–638.
116. Jensen BL, Friis UG, Hansen PB *et al*. Voltage-dependent calcium channels in the renal microcirculation. *Nephrol Dial Transplant* 2004; **19**: 1368–1373.
117. Kurtz A, Penner R. Angiotensin II induces oscillations of intracellular calcium and blocks anomalous inward rectifying potassium current in mouse renal juxtaglomerular cells. *Proc Natl Acad Sci USA* 1989; **86**: 3423–3427.
118. Taugner R, Schiller A, Kaissling B *et al*. Gap junctional coupling between the JGA and the glomerular tuft. *Cell Tissue Res* 1978; **237**: 279–285.
119. Krattinger N, Capponi A, Mazzolai L *et al*. Connexin40 regulates renin production and blood pressure. *Kidney Int* 2007; **72**: 814–822.
120. Haefliger J-A, Krattinger N, Martin D *et al*. Connexin43-dependent mechanism modulates renin secretion and hypertension. *J Clin Invest* 2006; **116**: 403–413.
121. Rackauskas M, Kreuzberg MM, Pranevicius M *et al*. Gating properties of heterotypic gap junction channels formed of connexin 40, 43, and 45. *Biophys J* 2007; **92**: 1952–1965.
122. Haefliger JA, Meda P, Formenton A *et al*. Aortic connexin 43 is decreased during hypertension induced by inhibition of nitric oxide synthase. *Arterioscler Thromb Vasc Biol* 1999; **19**: 1615–1622.
123. Hoffmann A, Gloe T, Pohl Z *et al*. Nitric oxide enhances *de novo* formation of endothelial gap junctions. *Cardiovasc Res* 2003; **60**: 421–430.
124. Zhang JH, Kawashima S, Yokoyama M *et al*. Increased eNOS accounts for changes in connexin expression in renal arterioles during diabetes. *Anat Rec A Discov Mol Cell Evol Biol* 2006; **288**: 1000–1008.
125. Zhang X, Chen X, Wu D *et al*. Downregulation of connexin 43 expression by high glucose induces senescence in glomerular mesangial cells. *J Am Soc Nephrol* 2006; **17**: 1532–1542.
126. Yao J, Hiramatsu N, Zhu Y *et al*. Nitric oxide-mediated regulation of connexin 43 expression and gap junctional intercellular communication in mesangial cells. *J Am Soc Nephrol* 2005; **16**: 58–67.
127. Haefliger JA, Castillo E, Waerber G *et al*. Hypertension increases connexin 43 in a tissue specific manner. *Circulation* 1997; **95**: 1007–1014.
128. Firouzi M, Kok B, Spiering W *et al*. Polymorphisms in human connexin 40 gene promoter are associated with increased risk of hypertension in men. *J Hypertens* 2005; **24**: 325–330.